

Calc 12 - Chp 8.1

Note Title

2013-09-23

L'Hopital's Rule - please learn on your own.

Tips: Read Theorem 1
" Examples 1 & 2
" 8.1 Intro
" Theorem 1
" Theorem 2
" Example 3
" Theorem 2.
" Examples 4-6

Work thru Example 7, make sure you understand each step.

HW: pp. 423-424: 1-7 odds, 15-27 odds

If you have trouble, read example 7 again.

Work thru Examples 8-10.

HW: p. 424: 33-39 odds.



Calc 12 - Chp 8.2

Note Title

2013-09-23

Relative Rates of Growth

Definitions: for $f(x)$ & $g(x)$ positive & x sufficiently large

1. f grows faster than g (implies g grows slower than f) as $x \rightarrow \infty$ if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty \quad \text{or} \quad \lim_{x \rightarrow \infty} \frac{g(x)}{f(x)} = 0$$

2. f and g grow at the same rate as $x \rightarrow \infty$ if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L, \quad L \neq 0 \text{ and finite}$$

You should notice that L'Hopital's Rule might come handy for this section.

eg) Show that e^x grows faster than x^2

eg) Show that x grows faster than $\ln x$.

Transitivity of Growth Rates.

If f grows at the same rate as g as $x \rightarrow \infty$ and g grows at the same rate as h as $x \rightarrow \infty$, then f grows at the same rate as h .

eg) Show $\sqrt{x^3 - 5}$ and $(3\sqrt{x} + 1)^3$ have the same growth rate.

Definition: f of Smaller Order than g , if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0, \text{ then we write } f = o(g)$$

and say "f is little-oh of g"

Definition: f of at Most the Order of g , if

$$\frac{f(x)}{g(x)} \leq M, \text{ then we write } f = O(g)$$

and say "f is big-oh of g"

eg) Show $4 \ln x = o(x^2)$

eg) Show $2x + \cos x = O(x)$

HW: p. 431: 1-23 odds (please show 20)

Calc 12 - Chp 8.3 Part 1

Note Title

2013-09-23

Improper Integrals

$$\int_0^{\infty} e^{-x/3} dx =$$

As you know, we can't just substitute ∞ .

Definitions: As long as $f(x)$ is continuous on the interval

$$1. \int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

$$2. \int_{-\infty}^b f(x) dx = \lim_{a \rightarrow -\infty} \int_a^b f(x) dx$$

$$3. \int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx, \text{ where } c \in \mathbb{R}$$

Note: These definitions do not imply that the improper integral has a finite value. If the limit equals $\pm\infty$, we say the integral diverges. If the limit has a finite value, we say it converges.

Continuing previous eq)

$$\int_0^{\infty} e^{-x/3} dx =$$

eg) Evaluate $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$

Use def 3, let $c=0$.

eg) Evaluate $\int_1^{\infty} \frac{2}{x} dx$

Theorem: $\int_1^{\infty} \frac{1}{x^p} dx$ is convergent if $p > 1$ else divergent.

Another improper integral happens when the interval ends at a vertical asymptote.

Definitions:

If f is continuous on $[a, b)$, then

$$\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx$$

If f is continuous on $(a, b]$, then

$$\int_a^b f(x) dx = \lim_{c \rightarrow a^+} \int_c^b f(x) dx$$

If f is continuous on $[a, c) \cup (c, b]$, then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

Again, we must determine if the improper integral converges or diverges.

eg) Find $\int_2^5 \frac{1}{\sqrt{x-2}} dx$

V.A. @ $x=2$

eg) Find $\int_0^3 \frac{1}{x-1} dx$

V.A. @ $x=1$

HW: p. 442: 1-21 odds.
(Please solve 14 in person)

Calc 12 - Chp 8.3 Part 2

Note Title

2013-09-23

Improper Integrals - Convergence Tests.

Some integrals are difficult to find, so to test for convergence, we look for an known integral to compare.

Theorem 3 - Direct Comparison Test.

Let f and g be continuous on $[a, \infty)$ with $0 \leq f(x) \leq g(x)$ for all $x \geq a$, then

1. $\int_a^{\infty} f(x) dx$ converges if $\int_a^{\infty} g(x) dx$ converges.
2. $\int_a^{\infty} g(x) dx$ diverges if $\int_a^{\infty} f(x) dx$ diverges.

This is like a one-sided Sandwich Theorem.

eg) Is $\int_0^{\infty} e^{-x^2} dx$ convergent?

Substitution won't work:

$$u = -x^2$$
$$du = \underbrace{-2x dx}_{\text{not there.}}$$

$\int e^{-x} dx$ is easily integrable.

eg) Is $\int_1^{\infty} \frac{1+e^{-x}}{x} dx$ convergent?

We notice that $\frac{1+e^{-x}}{x} > \frac{1}{x}$,

Theorem 4: Limit Comparison Test

If f and g are continuous on $[a, \infty)$, $f, g > 0$ and

$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L$, $L \in \mathbb{R}$, then

$\int_a^{\infty} f(x) dx$ and $\int_a^{\infty} g(x) dx$ both converge or both diverge

eg) Does $\int_1^{\infty} \frac{1}{x+x^2} dx$ converge?

Try comparing with $\int_1^{\infty} \frac{1}{x^2} dx$

HW: p. 442: 27-43 odds
(Please solve 38 in person)

Calc 12 - Chp 8.4

Note Title

2013-09-23

Partial Fractions and Integral Tables.

Most polynomial denominators are difficult to integrate directly, so partial fractions is another tool for our toolbox. We must solve a system of linear equations when using partial fractions.

Method of Partial Fractions ($f(x)/g(x)$ Proper)

1. Let $x - r$ be a linear factor of $g(x)$. Suppose $(x - r)^m$ is the highest power of $x - r$ that divides $g(x)$. Then, to this factor, assign the sum of the m partial fractions:

$$\frac{A_1}{x - r} + \frac{A_2}{(x - r)^2} + \dots + \frac{A_m}{(x - r)^m}.$$

Do this for each distinct linear factor of $g(x)$.

2. Let $x^2 + px + q$ be a quadratic factor of $g(x)$. Suppose $(x^2 + px + q)^n$ is the highest power of this factor that divides $g(x)$. Then, to this factor, assign the sum of the n partial fractions:

$$\frac{B_1x + C_1}{x^2 + px + q} + \frac{B_2x + C_2}{(x^2 + px + q)^2} + \dots + \frac{B_nx + C_n}{(x^2 + px + q)^n}.$$

Do this for each distinct quadratic factor of $g(x)$ that cannot be factored into linear factors with real coefficients.

3. Set the original fraction $f(x)/g(x)$ equal to the sum of all these partial fractions. Clear the resulting equation of fractions and arrange the terms in decreasing powers of x .
4. Equate the coefficients of corresponding powers of x and solve the resulting equations for the undetermined coefficients.

p. 445

Simple eg) $\int \frac{1}{x^2 - a^2} dx$ *

Med

eg)

$$\int \frac{x^2 + 2x - 1}{2x^3 + 3x^2 - 2x} dx$$

Hard eg)

$$\int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx$$

$$\begin{array}{r} x^3 - x^2 - x + 1 \overline{) x^4 + 0x^3 - 2x^2 + 4x + 1} \\ \underline{x^4 - x^3 - x^2 + x} \\ 0 x^3 - x^2 + 3x + 1 \\ \underline{x^3 - x^2 - x + 1} \\ \phantom{0 x^3 - x^2 +} 4x \end{array}$$

HW: p. 452: 1-13 odds, 17, 21, 25
(Please show 18 in person)